

Supplementary  
Table 1

**SITES OF METASTASES IN PATIENTS IN CELL TRANSFER PROTOCOLS**

<b>Category</b>	<b>No Gy</b>	<b>2Gy</b>	<b>12Gy</b>	<b>Total*</b>
Number of patients	43	25	25	93
Sites of Disease %				
Lung	40%	64%	24%	42%
Liver	19%	36%	16%	23%
Bone	9%	24%	4%	12%
Brain	26%	16%	0%	16%
Intraperitoneal	44%	56%	32%	44%
Peripheral LN	60%	44%	48%	53%
Skin/s.c.	79%	48%	20%	55%
Adrenal	12%	20%	24%	17%
Thoracic/mediastinal	19%	24%	16%	19%
Other	30%	20%	24%	26%
Skin and periph LN only	16%	12%	32%	19%
Lung only	0%	0%	4%	1%
Median number sites/patient	3	4	2	3
TNM Stage M1c	74%	92%	68%	77%
M1b/c	88%	96%	72%	86%

**STAGES OF METASTATIC DISEASE**

	CR*	PR*	NR*	Total
	(number of patients)			
Not BI: M1a**	1	1	3	5
1b	2	3	1	6
1c	2	12	18	32
2 Gy TBI: M1a	0	0	1	1
1b	0	1	0	1
1c	5	7	11	23
12 Gy TBI: M1a	2	3	2	7
1b	1	0	0	1
1c	7	5	5	17

\*CR, complete response; PR partial response; NR no response

\*\*M1a, skin, subcutaneous or nodal metastases; M1b lung; M1c all other visceral sites or elevated lactic dehydrogenase